



PENINSULA PULMONARY FUNCTION LABORATORY

ACCREDITED LABORATORY (TSANZ)

ABN: 79 103 204 164

Medical Director

Dr Keith Burgess

MB BS MSc Ph.D FRACP FRCPC

Clinical Associate Professor Dept of Medicine University Sydney

Frenchs Forest Laboratory

Building 1 Level 2 Unit 2

49 Frenchs Forest Road

FRENCHS FOREST 2086

Phone: 99754911

Fax: 99754622

PATIENT DETAILS

Name: _____

Address: _____

Date of Birth: _____

Mater Clinic Laboratory

Suite 1.05,

3-9 Gillies Street

CROWS NEST 2060

Smoking: YES

NO

Pack years?

Hb, date taken:

CLINICAL HISTORY

TESTS REQUIRED: please tick box

- Full pulmonary function (flow volume loop, with bronchodilator, lung volumes, gas transfer capacity)
- Flow volume loop / Spirometry, includes bronchodilator
- Airway Resistance & Conductance (RAW)
- Maximum Voluntary Ventilation (MVV)
- MIPS & MEPS (mouth pressures)
- Sniff Nasal Inspiratory Pressure (SNIP)
- Home Overnight Oximetry *
- High Altitude Simulation Test *

* Available only at Frenchs Forest

Tests for Asthma / Allergy

- Histamine Challenge
- Saline Challenge *
- Mannitol Challenge
- Nasal Resistance
- Skin Test for Allergies

Referring Doctor:

Provider number:

Address:

Date: / /

Pulmonary Function, Spirometry & Flow Volume Loops

- No short acting bronchodilators FOUR HOURS prior to test such as Ventolin, Bricanyl and Atrovent.
- No long acting bronchodilators 12hrs prior to test such as Serevent, Seretide, Oxis and Symbicort.
- No Onbrez, Spiriva, Seebri, Bretaris, Ultibro, Breo, Anoro, Incruse or Singulair for 48 hours prior to testing.
- No smoking for two hours prior to testing (ideally 24 hours).

Histamine / Saline / Mannitol Challenge

- No short acting bronchodilators 12 hours prior to test such as Ventolin, Bricanyl and Atrovent.
- No Serevent, Oxis, Onbrez, Spiriva, Seebri, Bretaris, Ultibro, Breo, Anoro or Incruse for at least 48 hours prior to testing
- For asthma diagnosis ideally no Flixotide, Intal, Pulmicort, Seretide, Symbicort , Qvar, Alvesco or Tilade for two weeks before test. (Although tests performed on therapy to assess effectiveness).
- No antihistamines, theophylline or Singulair tablets for at least 48hrs prior to testing.
- Oral Prednisolone, etc (cortisone)

(NB: Please discuss with prescribing doctor before withholding oral cortisone medication – Test not valid if taking this medication)

Nasal Resistance

- No vasoconstrictors 4hrs prior to test such as Drixine, Dimetapp.

Skin Tests

- Ideally should not be done if patient is on Prednisone \geq 5mg/day No antihistamines for at least 48hrs prior to testing.

HAST

- Patients should be stable at time of testing. May take all usual medications (unless combined with other tests eg. provocation test for asthma).

**For Directions, please see Website for Map –
www.peninsularespiratory.com.au**